

# Spatial Abilities of Expert Clinical Anatomists: Comparison of Abilities Between Novices, Intermediates, and Experts in Anatomy

Ruth Fernandez,<sup>1</sup> Itiel E. Dror,<sup>2,3</sup> Claire Smith<sup>1\*</sup>

<sup>1</sup>Centre for Learning Anatomical Sciences, Faculty of Medicine, University of Southampton, Southampton, Hampshire, United Kingdom

<sup>2</sup>Institute of Cognitive Neuroscience, University College London, London, England

<sup>3</sup>Cognitive Consultants International Ltd., Southampton, Hampshire, England

Spatial ability has been found to be a good predictor of success in learning anatomy. However, little research has explored whether spatial ability can be improved through anatomy education and experience. This study had two aims: (1) to determine if spatial ability is a learned or inherent facet in learning anatomy and (2) to ascertain if there is any difference in spatial ability between experts and novices in anatomy. Fifty participants were identified: 10 controls, 10 novices, 10 intermediates, and 20 experts. Participants completed four computerized spatial ability tasks, a visual mental rotation task, categorical spatial judgment task, metric spatial task, and an image-scanning task. The findings revealed that experts ( $P = 0.007$ ) and intermediates ( $P = 0.016$ ) were better in the metric spatial task than novices in terms of making more correct spatial judgments. Experts ( $P = 0.033$ ), intermediates ( $P = 0.003$ ), and novices ( $P = 0.004$ ) were better in the categorical spatial task than controls in terms of speed of responses. These results suggest that certain spatial cognitive abilities are especially important and characteristic of work needed in clinical anatomy, and that education and experience contribute to further development of these abilities. *Anat Sci Educ* 00: 000–000. © 2011 American Association of Anatomists.

*Key words:* anatomical sciences; medical education; anatomy education; computers in anatomy education; spatial abilities; mental rotation

## INTRODUCTION

It has been suggested in recent literature that changes in medical training over the years have increasingly neglected anatomy (Ellis, 2002) and that this has, in part, been the cause for the recent increase of medicolegal claims against surgeons in the United Kingdom (Goodwin, 2000). Moreover, new

styles in anatomy teaching have taken much criticism (Kaufman, 1997; Hanna and Tang, 2005), especially teaching that focuses on computer and textbook-based learning as opposed to more traditional hands-on dissection techniques (Amadio, 1996; Cahill and Leonard, 1997; Ellis, 2001; von Lüdinghausen, 2001; Korf et al., 2008; Wood et al., 2010). A focus on anatomy teaching during medical training is necessary to identify where practical improvements can be made. Spatial ability has been found to be a good predictor of students' success in learning anatomy and examination performance (Garg et al., 2001). It has also been suggested that spatial ability might be even more important than the type of educational materials that are studied (Garg et al., 2001). Spatial ability has also been related to clinical performance. Wanzel et al. (2003) suggested that through experience, surgical performance increases regardless of individual spatial ability (or manual dexterity) making the case that inherent spatial ability becomes less important as experience takes over.

\*Correspondence to: Dr. Claire Smith, Centre for Learning Anatomical Sciences, Faculty of Medicine, University of Southampton, South Academic Block, Southampton General Hospital, Tremona Road, Southampton, Hampshire SO16 6YD, United Kingdom. E-mail: c.f.smith@southampton.ac.uk

Received 19 August 2010; Revised 17 November 2010; Accepted 1 December 2010.

Published online in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/ase.196

Despite its effect on performance, both academically and clinically, spatial learning is poorly understood, and there has been very little research into the various components of spatial abilities and their implications to teaching anatomy.

Studies related to spatial ability and anatomy experience are limited; however, it is evident that experience can improve spatial ability to some extent. Kioumourtzoglou et al. (1998) found that water polo players have significantly better scores than novices on decision making, visual reaction time, and spatial orientation. Furthermore, Dror et al. (1993) found that pilots judged metric spatial relations better than nonpilots, and pilots mentally rotated objects better than nonpilots, again providing evidence for links between spatial ability and experience. With the current concerns over lamentable anatomy knowledge gained at medical school and rising litigation linked to a deficit of anatomical knowledge (Ellis, 2002; Older, 2004), understanding spatial ability with the view to improving anatomy teaching would be valuable.

The aim of this study was to examine spatial ability in experts and novices in anatomy. We used the four experimental paradigms developed by Dror et al. (1993) to objectively test and quantify various components of spatial ability of people with different experience and training in medical anatomy, from total novices to experts.

## MATERIALS AND METHODS

### Study Design

This comparative study gained favorable ethics approval from the University of Southampton, School of Medicine (SOMSEC030.09).

### Participants

Random sampling was used to select 10 controls. Ten novices (first-year medical students), 10 intermediates (fourth- and fifth-year medical students), and 20 experts. Experts for the purpose of this study were defined as university lecturers who had more than five years' experience of teaching anatomy. The demographics of the control population included an age range of 20–50 years, an academic qualification not higher than GCSE and occupations that were not related to spatial ability, e.g., anatomy technicians, architects, and pilots. There were an equal number of males and females for each category as a gender difference may exist (e.g., Voyer et al., 2000; Peters et al., 2007). Informed consent was gained from all participants.

### Methods of Measurement

The method has been designed specifically to explore four components of spatial abilities related to anatomy, components of which had been used previously to study military fighter pilots (Dror et al., 1993). These were administered via computerized experiments: a visual mental rotation task, a scanning task, a categorical spatial relation task, and a metric spatial relation task. The tasks were administered and counterbalanced across participants. Participants first had practice trials to familiarize themselves with the experimental setup and tasks. For the practice trials, participants received feedback. It was confirmed that none of the participants had previous experience with these computerized experiments.

**Rotation task.** The rotation task was selected as it mimics the need for orientation in anatomy. This type of spatial appreciation may take place clinically, for example, when orientating anatomical structures on a CT image.

Participants were presented with two consecutive black and white drawings. The first drawing was always presented upright, whereas the second drawing was either identical or differed slightly from the first presentation (e.g., an additional line was present or absent, or a shape was changed). The second drawing was presented at 0°, 35°, 70°, or 105°. The participants were required to judge whether the two drawings were the same, regardless of orientation. This task consisted of 48 trials. The trials were presented in a fixed pseudo-random order with no more than three consecutive “yes” or “no” trials, no more than three consecutive trials with the second stimuli in the same orientation, and no more than three consecutive trials with the same objects. All participants received the same order of presentation.

**Scanning task.** This task was selected as it tests the ability to scan images and recall positions of objects without the relevant stimuli. In clinical practice, this is used, for example, for image-guided procedures such as angioplasty.

Participants were presented with a circle consisting of 16 segments. Three of these segments were black and the others white. All the segments then turned white and an arrow appeared in the center of the circle. Participants were required to judge whether or not the segment to which the arrow was pointing was previously black. This task consisted of 48 trials in which the arrow pointed to a segment which was previously black in one half of the trials, and the other half pointed to a segment which was previously white. The trials were presented in a fixed pseudo-random order, with no more than three consecutive “yes” or “no” trials, no more than three consecutive trials with the arrow pointing to the same segment, and no more than three consecutive trials with the arrow pointing from the same distance. All participants received the same order of presentation.

**Categorical task.** This task was selected as it tests the ability to judge the categorical relation of one object to another. This is used clinically, for example, when surface anatomical landmarks are used to find underlying structures.

Participants were presented with a dot located above or below a bar and asked to judge whether the dot was above the bar. The dot could appear at one of four distances away from the bar. The bar could appear in one of three locations; centrally and slightly above and below central. Bar-dot stimuli were presented in a fixed pseudo-random order, with no more than three consecutive “above” or “below” trials and no more than three consecutive trials with the dot being a certain distance away from the bar. All participants received the stimuli in the same order.

**Metric task.** The metric task was selected as it tests the ability to judge specific distance. This is used, for example, when surgeons need to appreciate depth of fascial layers or when clinicians take blood.

Participants were presented with a dot located above or below a bar at different distances. For this task, participants were required to estimate the distance between the dot and the bar. Exactly the same stimuli were used in this task as in the categorical task; however, participants were required to make a different type of spatial judgment. Participants made their response by typing in their estimation (in cm) using the number pad on the keyboard with their dominant hand.

**Table 1.**

Mean Number of Correct Answers

Task	Participants	N	Mean	95% Confidence interval for mean		Minimum	Maximum
				Lower bound	Upper bound		
Metric	Intermediates	10	36.90	29.87	43.93	17	47
	Experts	20	36.55	32.33	40.77	12	47
	Novices	10	25.30	15.17	35.43	3	43
	Control	10	33.40	26.92	39.88	19	45
	Total	50	33.74	30.61	36.87	3	47
Scanning	Intermediates	10	43.60	40.60	46.60	33	47
	Experts	20	41.55	39.42	43.68	30	48
	Novices	10	43.30	41.11	45.49	37	47
	Control	10	40.60	37.75	43.45	35	48
	Total	50	42.12	40.94	43.30	30	48
Categorical	Intermediates	10	47.00	45.99	48.01	44	48
	Experts	20	47.40	46.87	47.93	44	48
	Novices	10	46.20	43.77	48.63	37	48
	Control	10	47.40	46.71	48.09	45	48
	Total	50	47.08	46.56	47.60	37	48
Rotation	Intermediates	10	39.70	36.98	42.42	31	44
	Experts	20	38.55	36.53	40.57	31	47
	Novices	10	37.80	34.62	40.98	31	43
	Control	10	38.10	34.19	42.01	31	46
	Total	50	38.54	37.29	39.79	31	47

## Statistical Analysis

The numbers of correct answers and the mean response times were subjected to analysis of variance, comparing performance among the control, novice, intermediate, and expert groups on each of the tasks (using a 95% confidence interval). A post hoc test was performed looking at least significant difference (LSD) and pair-wise comparisons between the groups. A Bonferroni correction was performed to counter the effects of multiple testing.

## RESULTS

### Number of Correct Answers

T1 The metric task revealed that experts and intermediates were performing better than the other groups (see Table 1). Results from parametric statistics performed on this data confirmed

the statistical differences between groups;  $P = 0.04$  (see Table 2). In all the other tasks (scanning, categorical, and rotation), no significant difference was found (see Table 2). A post hoc test was performed on the data from the metric task looking at LSD and pair-wise comparisons between the groups (see Table 3). Significant difference was found T2 T3 between the intermediate and novice groups ( $P = 0.016$ ) and expert and novice groups ( $P = 0.007$ ). The intermediates and experts scored a significantly higher number of correct answers than the novices. A Bonferroni correction test was completed and showed that even when accounting for multiple testing, the mean difference between experts and novices is significant,  $P = 0.045$  (see Table 3).

### Response Times

Significant differences were found between groups for the scanning task,  $P = 0.05$ , and the categorical spatial task,  $P = 0.10$  (see Table 4). A post hoc test was performed for a pair- T4

**Table 2.**

Mean Number of Correct Answers—Overall ANOVA

Task	Comparison	df	<i>P</i> value
Metric	Between groups	3	0.040
	Within groups	46	
	Total	49	
Scanning	Between groups	3	0.290
	Within groups	46	
	Total	49	
Categorical	Between groups	3	0.368
	Within groups	46	
	Total	49	
Rotation	Between groups	3	0.793
	Within groups	46	
	Total	49	

df, degree of freedom.

wise comparison between groups (see Table 5). Average response time was found to be significantly faster in the scanning task in the intermediates than experts ( $P = 0.027$ ) and controls ( $P = 0.033$ ). Intermediates ( $P = 0.003$ ), experts ( $P = 0.033$ ), and novices ( $P = 0.004$ ) response times were all found to be significantly faster than controls in the categorical spatial task.

## DISCUSSION

For the metric spatial relations task, both the expert and intermediate groups outperformed the novice group, the experts and the intermediates scoring significantly higher than the novices. These results suggest that ability to judge distance may have improved through experience.

Although no significant difference was found in the categorical spatial relations task between groups when examining the number of correct answers, differences were found in response times. Novices, intermediates, and experts all responded significantly faster than the control group. These results suggest that response times in ability to assess spatial relations may have also been improved through experience. The image-scanning task showed no significant differences between groups when looking at the number of correct answers; however, differences were found in response times. Intermediates scored significantly higher than both the control and expert groups. These results suggest that response times in ability to recall positions were better in the student groups. One possible explanation for these results may be

**Table 3.**

Mean Number of Correct Answers: Metric Task: Post Hoc Test: Least Significant Difference (LSD) and Bonferroni Correction (BC)

	(I) Group	(J) Group	Mean difference (I – J)	<i>P</i> value	95% Confidence interval	
					Lower bound	Upper bound
LSD	Intermediates	Experts	0.350	0.931	-7.74	8.44
		Novices	11.600 <sup>a</sup>	0.016	2.25	20.95
		Control	3.500	0.455	-5.85	12.85
Experts	Novices	Control	11.250 <sup>a</sup>	0.007	3.16	19.34
		Control	3.150	0.437	-4.94	11.24
Control	Novices	8.100	0.088	-1.25	17.45	
BC	Intermediates	Experts	0.350	1.000	-10.74	11.44
		Novices	11.600	0.097	-1.20	24.40
		Control	3.500	1.000	-9.30	16.30
Experts	Novices	Control	11.250 <sup>a</sup>	0.045	0.16	22.34
		Control	3.150	1.000	-7.94	14.24
Control	Novices	8.100	0.526	-4.70	20.90	

Comparing (I) group, the control, to (J) group, the comparator (representing intermediates, experts, novices, and control groups, respectively), to see if there is a significant difference in results between these groups.

<sup>a</sup>Significant at  $P$  value ( $<0.05$ ) showing (J) group to be inferior to (I) group.

**Table 4.**

## Average Response Times

Average response time in tasks	Comparison	df	P value
Metric	Between groups	3	0.546
	Within groups	46	
	Total	49	
Scanning	Between groups	3	0.050
	Within groups	46	
	Total	49	
Categorical	Between groups	3	0.010
	Within groups	46	
	Total	49	
Rotation	Between groups	3	0.801
	Within groups	46	
	Total	49	

df, degree of freedom

that student participants are more familiar with computer-based programs than the expert group (because of a generation difference), and, therefore, the student groups were quicker at responding on these computer-based programs. For mental rotation, no significant differences were found between any of the groups in either the number of correct answers or response times. These results suggest that the ability to mentally rotate had not been improved through experience.

In most of the tasks, it was found that expert groups responded faster overall than nonexpert groups (with the exception of image scanning). We found that experts had better ability to judge metric spatial relations. However, in contrast, there was no evidence that the more expert groups have higher ability to scan visual mental images or mentally rotate objects. Thus, there is evidence that experts have selective advantages, not overall superior performance. Reasons for selective advantages in the more experienced groups may be focused on the variable plasticity of different regions of the brain responsible for difference aspects of spatial ability. Evidence suggests that some processes in the brain are more plastic and thus susceptible to change, whereas other processes are less plastic, and one possible reason for such differences is that some processes rely on more primitive, hard-wired brain structures than others do (DeFelipe, 2006).

The metric spatial relations task requires the participant to make precise distance judgments. Such processing relies on accurately making small spatial distinctions, which involves the parietal lobes, particularly right parietal lobe structures (Hellige and Michimata, 1989; Kosslyn et al., 1989). However, no difference was found in the mental rotation task where such processing relies on a set of complex computa-

tions that involve parietal and frontal lobe structures (Deutsch et al., 1988).

All groups scanned images at comparable rates finding no significant differences between levels of expertise; image scanning is thought to involve the middle temporal area of the brain, possibly suggesting that image scanning is a less adaptable brain process (Allman et al., 1985).

Overall, the more experienced groups judged metric spatial relations better than the novice group. Other studies have also demonstrated this same finding. For example, Dror et al. (1993) showed that ability to judge metric spatial relations is learnt through experience. Both the expert and intermediate groups outperformed the novices, and this result dovetails well with other research, which suggest that inherent spatial ability becomes less important as experience takes over (Wanzel et al., 2003). It is clear from previous research that spatial ability is a reliable predictor of success in learning anatomy (Rochford, 1985; Garg et al., 2001; Guillot et al., 2006).

This is all a prelude to focusing anatomy teaching on developing those spatial skills that are susceptible to change. If spatial abilities are adaptable in individuals, then improving 3D anatomy teaching, techniques, and materials is paramount in advancing anatomy learning amongst medical students. The review of current teaching techniques enters into the ongoing debate over the advantages and disadvantages of using human cadavers for teaching. The ability to observe the form of 3D structures and the spatial relationship between them are some of the primary advantages of learning anatomy by using human cadavers (Crisp, 1989; Hill and Anderson, 1991; Pabst, 1993; Marks, 1996; Wood et al., 2010).

Evidence also suggests that computer-based teaching materials are associated with a better understanding of spatial anatomy (Silén et al., 2008; Petersson et al., 2009) and improved learning (Lynch et al., 2001; St Aubin, 2001; McNulty et al., 2004, 2009; Sugand et al., 2010). Furthermore, they have proved to be well received by students (Nieder et al., 2000; McNulty et al., 2009). Whether human cadavers have a select advantage over computer-based material is a matter of ongoing debate; however, it is clear that both techniques use spatial ability and could have the potential to improve individual spatial abilities that are susceptible to change. Moreover, measuring the spatial abilities that are not susceptible to change could be used as criteria for screening and selecting medical professionals of which anatomical spatial ability is most relevant, for example, surgeons.

Although it might be thought that interest in applied anatomy may be driven by innate higher spatial ability, evidence suggests that individual interest is actually governed by perceived training needs (Langlois et al., 2009). Therefore, medical trainees may not necessarily be choosing medical professions that compliment their innate skill strengths further highlighting the importance of attempting to introduce more techniques and teaching methods to improve spatial ability where possible.

The authors recognize that the study had some limitations. First, the experimental design used two-dimensional (2D), not three-dimensional (3D) pictures. Anatomical spatial appreciation involves 3D and 2D visualization. As this was the first study of its kind, 2D testing was appropriate and has established a baseline; however a later study could build on the data to include 3D testing. However, the cognitive literature suggests that 2D and 3D image rotations are very similar

**Table 5.**

Average Response Times: Post Hoc Test

Dependent variable	(J) Group	(I) Group	P value	95% Confidence interval	
				Lower bound	Upper bound
Metric average response time	Experts	Intermediates	0.167	-131.6036	737.5856
		Novices	0.389	-246.6606	622.5286
		Control	0.597	-319.5736	549.6156
	Novices	Intermediates	0.647	-386.7696	616.8836
		Control	0.455	-313.8566	689.7966
		Novices	0.771	-428.9136	574.7396
Scanning average response time	Experts	Intermediates	0.027	76.0099	1219.9681
		Novices	0.077	-57.5881	1086.3701
		Novices	0.686	-526.8666	794.0626
	Control	Intermediates	0.033	59.7944	1380.7236
		Experts	0.800	-499.7091	644.2491
		Novices	0.080	-73.8036	1247.1256
Categorical average response time	Experts	Intermediates	0.149	-32.1764	205.4974
		Novices	0.206	-43.1644	194.5094
		Novices	0.873	-126.2330	148.2090
	Control	Intermediates	0.003	79.0660	353.5080
		Experts	0.033	10.7896	248.4634
		Novices	0.004	68.0780	342.5200
Rotation average response time	Intermediates	Experts	0.841	-710.5326	868.9756
		Novices	0.513	-613.4274	1210.4314
		Experts	0.579	-570.4736	1009.0346
	Control	Intermediates	0.752	-767.8329	1056.0259
		Experts	0.572	-566.4361	1013.0721
		Novices	0.334	-469.3309	1354.5279

Comparing (I) group, the control, to (J) group, the comparator (representing intermediates, experts, novices, and control groups, respectively), to see if there is a significant difference in results between these groups. A significant *P* value (<0.05) showing (J) group to be inferior to (I) group.

from a cognitive perspective. Genetic, hormonal, and neurological factors were not controlled and have been found to potentially affect individual spatial ability (McGee, 1979). Environmental factors, such as vocational activities involving spatial intelligence, were also not controlled. Although participants' ages were noted, this factor was not taken into account when analyzing results but has also been shown to

potentially influence spatial ability (Salthouse and Mitchell, 1990; Salthouse et al., 1990). It is possible that self-selection bias meant that the participants who volunteered to take part in the study were those who believe that they have good spatial ability and thus introduced self-selection bias. Furthermore, the medical students used for this study were all from the University of Southampton, whereas the anatomy

experts had studied in a range of universities. As the Southampton students had received all the same anatomy teaching, it might be that this university may teach spatially relevant anatomy differently compared with other universities, and so these students may score a higher or lower average on the tasks. For the categorical and scanning tasks, the majority of scores are at or near the maximum possible for the test (48), suggesting a possible ceiling effect. This may have limited the ability to measure the different groups by creating values near the ceiling limit. This may have reduced variance, decreasing the sensitivity of the experiment and, therefore, does not determine if the average of one group is significantly different from the average of another group. The control group involved a range of ages 20–50 years; in subsequent tests, it may be possible to improve age matching and have a control group for each participant group, e.g., nonmedical students and nonscience professors.

## CONCLUSION

This study is a step toward understanding spatial abilities required for conducting and understanding anatomy. Spatial ability is a good predictor of successful anatomy learning, and evidence from this study suggests that only certain aspects of spatial ability can be gained through experience. Current educational methods, such as dissection, and more recently 3D computer images, have been shown to produce a better understanding of spatial anatomy (Crisp, 1989; Hill and Anderson, 1991; Pabst, 1993; Marks, 1996; Silén et al., 2008; Petersson et al., 2009). In the light of the results from this study, it may be possible to develop “spatial ability” sessions, which at various points in the curriculum enable students to refine their spatial skills in order to enable more effective learning of anatomy. This would add to current methods of learning through specifically targeting certain components of spatial ability, as this is important in successful anatomy learning (Garg et al., 2001).

## NOTES ON CONTRIBUTORS

RUTH FERNANDEZ, B.Med., B.Sc., is a foundation Year 1 doctor in the Royal Hampshire County Hospital, Winchester, United Kingdom. She undertook part of the experimental work and performed data analysis.

ITIEL E. DROR, Ph.D., is a researcher in the Institute of Cognitive Neuroscience at the University College London, London, United Kingdom, and a principal consultant and researcher at the Cognitive Consultants International Ltd., Southampton, United Kingdom. He advised on the experimental design.

CLAIRE F. SMITH, Ph.D., is a senior teaching fellow in the Centre for Learning Anatomical Sciences, University of Southampton, School of Medicine, Southampton, United Kingdom. She initiated the project and undertook part of the experimental work.

## ACKNOWLEDGMENTS

The authors thank all participants who made this study possible. They also thank Scott Harris for his advice on the statistical analysis and the University of Southampton School of Psychology for designing the computer tasks.

## LITERATURE CITED

- Allman J, Miezin F, McGuinness E. 1985. Direction- and velocity-specific responses from beyond the classical receptive field in the middle temporal visual area (MT). *Perception* 14:105–126.
- Amadio PC. 1996. Reaffirming the importance of dissection. *Clin Anat* 9:136–137.
- Cahill DR, Leonard RJ. 1997. The role of computers and dissection in teaching anatomy: A comment. *Clin Anat* 10:140–141.
- Crisp AH. 1989. The relevance of anatomy and morbid anatomy for medical practice and hence for postgraduate and continuing medical education of doctors. *Postgrad Med J* 65:221–223.
- DeFelipe J. 2006. Brain plasticity and mental processes: Cajal again. *Nat Rev Neurosci* 7:811–817.
- Deutsch G, Bourbon WT, Papanicolaou AC, Eisenberg HM. 1988. Visuospatial tasks compared via activation of regional cerebral blood flow. *Neuropsychologia* 4:2051–2062.
- Dror IE, Kosslyn SM, Waag WL. 1993. Visual-spatial abilities of pilots. *J Appl Psychol* 78:763–773.
- Ellis H. 2001. Teaching in the dissecting room. *Clin Anat* 14:149–151.
- Ellis H. 2002. Medico-legal litigation and its links with surgical anatomy. *Surgery* 20:i–ii.
- Garg AX, Norman G, Sperotable L. 2001. How medical students learn spatial anatomy. *Lancet* 357:363–364.
- Goodwin H. 2000. Litigation and surgical practice in the UK. *Br J Surg* 87:977–979.
- Guillot A, Champely S, Batier C, Thiriet P, Collet C. 2006. Relationship between spatial abilities, mental rotation and functional anatomy learning. *Adv Health Sci Educ Theory Pract* 12:491–507.
- Hanna SJ, Tang T. 2005. Reduced undergraduate medical science teaching is detrimental for basic surgical training. *Clin Anat* 18:465–466.
- Hellige JB, Michimata C. 1989. Categorization versus distance: Hemispheric differences for processing spatial information. *Mem Cognit* 17:770–776.
- Hill RB, Anderson RE. 1991. The uses and value of autopsy in medical education as seen by pathology educators. *Acad Med* 66:97–100.
- Kaufman MH. 1997. Anatomy training for surgeons—A personal viewpoint. *J R Coll Surg Edinb* 42:215–216.
- Kioumourtzoglou E, Kourtessis T, Michalopoulou M, Derri V. 1998. Differences in several perceptual abilities between experts and novices in basketball, volleyball and water-polo. *Percept Mot Skills* 86:899–912.
- Korf HW, Wicht H, Snipes RL, Timmermans JP, Paulsen F, Rune G, Baumgart-Vogt E. 2008. The dissection course—Necessary and indispensable for teaching anatomy to medical students. *Ann Anat* 190:16–22.
- Kosslyn SM, Koenig O, Barrett A, Cave CB, Tang J, Gabrieli JD. 1989. Evidence for two types of spatial representations: Hemispheric specialization for categorical and coordinate relations. *J Exp Psychol Hum Percept Perform* 15:723–735.
- Langlois J, Wells GA, Lecourtois M, Bergeron G, Yetisir E, Martin M. 2009. Spatial abilities in an elective course of applied anatomy after a problem-based learning curriculum. *Anat Sci Educ* 2:107–112.
- Lynch TG, Steele DJ, Johnson Palensky JE, Lacy NL, Duffy SW. 2001. Learning preferences, computer attitudes, and test performance with computer-aided instruction. *Am J Surg* 181:368–371.
- Marks SC Jr. 1996. Information technology, medical education, and anatomy for the twenty-first century. *Clin Anat* 9:343–348.
- McGee MG. 1979. Human spatial abilities: Psychometric studies and environmental, genetic, hormonal, and neurological influences. *Psychol Bull* 86:889–918.
- McNulty JA, Halama J, Espiritu B. 2004. Evaluation of computer-aided instruction in the medical gross anatomy curriculum. *Clin Anat* 17:73–78.
- McNulty JA, Sonntag B, Sinacore JM. 2009. Evaluation of computer-aided instruction in a gross anatomy course: A six-year study. *Anat Sci Educ* 2:2–8.
- Nieder GL, Scott JN, Anderson MD. 2000. Using QuickTime virtual reality objects in computer-assisted instruction of gross anatomy: Yorick—The VR skull. *Clin Anat* 13:287–293.
- Older J. 2004. Anatomy: A must for teaching the next generation. *Surgeon* 2:79–90.
- Pabst R. 1993. Gross anatomy: An outdated subject or an essential part of a modern medical curriculum? Results of a questionnaire circulated to final-year medical students. *Anat Rec* 237:431–433.
- Peters M, Manning JT, Reimers S. 2007. The effects of sex, sexual orientation, and digit ratio (2D:4D) on mental rotation performance. *Arch Sex Behav* 36:251–260.
- Petersson H, Sinkvist D, Wang C, Smedby O. 2009. Web-based interactive 3D visualization as a tool for improved anatomy learning. *Anat Sci Educ* 2:61–68.
- Rochford K. 1985. Spatial learning disabilities and underachievement among university anatomy students. *Med Educ* 19:13–26.
- Salthouse TA, Mitchell DR. 1990. Effects of age and naturally occurring experience on spatial visualization performance. *Dev Psychol* 26:845–854.

- Salthouse TA, Babcock RL, Skovronek E, Mitchell DR, Palmon R. 1990. Age and experience effects in spatial visualization. *Dev Psychol* 26:128–136.
- Silén C, Wirell S, Kvist J, Nylander E, Smedby O. 2008. Advanced 3D visualization in student-centred medical education. *Med Teach* 30:e115–e124.
- St Aubin H. 2001. Implementing a virtual reality paradigm in human anatomy/physiology college curricula. *Stud Health Technol Inform* 81:475–478.
- Sugand K, Abrahams P, Khurana A. 2010. The anatomy of anatomy: A review for its modernization. *Anat Sci Educ* 3:83–93.
- von Lüdinghausen M. 2001. Replacing dissection courses with electronic substitutes. *Clin Anat* 14:93.
- Voyer D, Nolan C, Voyer S. 2000. The relation between experience and spatial performance in men and women. *Sex Roles* 43:891–915.
- Wanzel KR, Hamstra SJ, Caminiti MF, Anastakia DJ, Grober ED, Reznick RK. 2003. Visual-spatial ability correlates with efficiency of hand motion and successful surgical performance. *Surgery* 134:750–757.
- Wood A, Struthers K, Whiten S, Jackson D, Herrington CS. 2010. Introducing gross pathology to undergraduate medical students in the dissecting room. *Anat Sci Educ* 3:97–100.